



Philippine Society for Microbiology and Infectious Disease, Inc.

Subspecialty Society of the Philippine College of Physicians

Accredited Specialty Division of the Philippine Medical Association

Department of Science and Technology Accredited with Accreditation No. 14-F-05

Office address: 116 9th Avenue, Cubao, Quezon City 1109, Philippines • Tel. No.: (+632) 9126036;

Telefax No.: (+632) 9116986 • email address: psmid1970@gmail.com • website address: www.psmid.org

PSMID INTERIM GUIDE ON THE USE OF DENGUE VACCINE (Update, 6 December 2017)

National Officers 2017

Mari Rose A. De Los Reyes, MD, FPSMID
President

Mario M. Panaligan, MD, FPSMID
Vice President

Marissa M. Alejandria, MD, FPSMID
Secretary

Vegloure M. Maguinsay, MD, FPSMID
Treasurer

Henry F. Alavaren, MD, FPSMID
Business Manager

Councils

Janice C. Caoili, MD, FPSMID

Raul V. Destura, MD, FPSMID

Arthur Dessi E. Roman, MD, FPSMID

Elfleda A. Hernandez, MD, FPSMID

Larissa Lara Q. Torno, MD, FPSMID

Ellamae S. Divinagracia, MD, FPSMID

Immediate Past President

Marie Yvette C. Barez, MD, FPSMID

Chapter Presidents

Cebu

Belle M. Ranile, MD

Southern Mindanao

Larissa Lara Q. Torno, MD, FPSMID

Western Visayas

Lina C. Amsua, MD, FPSMID

Council of Advisers

Norma H. Abejar, MD, FPSMID

Rosario Angeles T. Alora, MD, FPSMID

Manolito L. Chua, MD, FPSMID

Remedios F. Coronel, MD, FPSMID

Salvacion R. Gatchalian, MD, FPSMID

Ludovico L. Jurao, Jr., MD, FPSMID

Evelina N. Lagamayo, MD, FPSMID

Mary Ann D. Lansang, MD, FPSMID

Julius A. Lecciones, MD, FPSMID

Ma. Cecilia S. Montalban, MD, FPSMID

Jaime C. Montoya, MD, FPSMID

Mediadora C. Saniel, MD, FPSMID

Rontgene M. Solante, MD, FPSMID

Enrique A. Tayag, MD, FPSMID

Thelma E. Tupasi, MD, FPSMID

BACKGROUND

Dengue is endemic in the Philippines and vaccination is an important component of a comprehensive dengue prevention and control strategy, in conjunction with vector control, surveillance and management of cases. In December 2015, the Philippines was the first Asian country to register and approve the use of the first dengue vaccine, Dengvaxia. A pooled analysis of the CYD 14/15 trials of the dengue vaccine showed higher efficacy rates for seropositive patients aged ≥ 9 years (81.9%, 95% CI 67.2, 90) compared to seronegative patients (52.5%, 95% CI 5.9-76.1).¹ According to the WHO Scientific Advisory Group of Experts on immunization, countries with dengue seroprevalence of $\geq 70\%$ should be targeted for vaccination.² PSMID subsequently released a position statement based on the best available scientific knowledge at that time.³ In its statement, PSMID noted safety signals in the clinical trials suggesting a potential risk of antibody-dependent enhancement (ADE) of breakthrough dengue infection when the vaccine is administered to seronegative individuals. The Society thus recommended additional studies, including long-term surveillance of vaccinated individuals, to validate this observation.

On the 29th November 2017, Sanofi Pasteur released news of additional safety and efficacy analyses of five years of clinical data on study participants in the vaccine trials in Asia and Latin America. The announcement stated continued protection against dengue among those who had prior dengue infection but recommended against vaccination of those without previous infection.⁴ According to the interim conditional recommendation of WHO released last 30 November 2017, dengue vaccine should only be restricted to individuals with prior exposure to dengue.



In addition, WHO noted that subjects without prior exposure to dengue (seronegative) had higher risk of severe dengue and hospitalizations should they contract the disease after vaccination.⁵ In the Dengvaxia trial, the criteria used in classifying dengue fever among the participants of was based on the grading system of WHO 1997 classification. The Independent Data Monitoring Committee classified participants as having severe dengue using the criteria: virologically confirmed dengue with at least one of the following: platelet count $<100,000^9/L$ and bleeding and plasma leakage; shock; bleeding requiring blood transfusion; encephalopathy; liver impairment; impaired kidney function; myocarditis, pericarditis or heart failure.⁶

PRACTICE

At present, dengue fever is classified into three types: dengue without warning signs, dengue with warning signs, and severe dengue. Based on the WHO 2009 Dengue classification, severe dengue is now defined as having clinical manifestations of dengue plus any of the following: severe plasma leakage leading to shock, or fluid accumulation with respiratory distress, severe bleeding and severe organ impairment (either as having AST or ALT > 1000 , presence of seizures or impaired consciousness, myocarditis or renal failure).⁷

Currently, there are no definitive tests that can determine the serostatus or past infection of dengue. In the dengue vaccine trial, dengue plaque reduction neutralization test (PNRT) was used to determine baseline serostatus of the participants.⁶ Unfortunately, this is currently not available for commercial use. In the additional analyses carried out for the cohort of vaccinees followed up for six years, an NS1 antibody assay was used to determine the serostatus at baseline. Antibody tests such as dengue IgG and IgM are meant to detect current or recent past dengue infection but is not accurate enough to determine past infection. These tests can also have cross-reaction; thus, they can also be positive if a person is infected with any flavivirus such as chikungunya.

INTERIM STATEMENT

Based on the above information, PSMID asserts that:

- Dengue vaccine is safe and effective among those with prior exposure to dengue. Current evidence does not support giving dengue vaccine to individuals without past dengue infection.
- Vaccines are not without risk; we support the right of each individual to full disclosure prior to consent. Disclosure should include the epidemiology of the disease, probable outcomes once disease is contracted and available treatments, and the short- and long-term risks and benefits of vaccination.



- Long-term follow-up and independent review of efficacy and safety of drugs are important aspects of patient safety. PSMID supports mechanisms and initiatives that would determine the seroprevalence of dengue in representative samples across geographic areas and age groups, the target population for vaccination, the safety and cost-effectiveness of current and future dengue vaccines.
- A national registry of individuals who received at least one dose of Dengvaxia should be created. These individuals should be monitored regularly for occurrence of signs and symptoms compatible with dengue for at least 6 years. They are encouraged to consult any physician once they have symptoms for proper management, work-up, and immediate notification to health authorities.
- For individuals who have yet to complete the 3 doses, they are encouraged to wait for further advisory from the Department of Health, pending review of the intention-to-treat analyses of the long-term efficacy of the vaccine.
- For those who intend to get vaccinated, a history of past dengue infection should be determined, and the option of serotesting prior to vaccination should be offered. Risks and benefits should be discussed prior to vaccination, as well as the risks of not being immunized.
- There is no simple solution to preventing dengue. Thus, to have a sustainable dengue control program, appropriate clinical management, laboratory surveillance and prevention efforts such as expansion of disease surveillance to include the communities and the private sector, vector control, and strengthening of epidemic response at all levels should always be emphasized.



REFERENCES

1. Hadinegoro SR, Arredondo-Garcia JL, Capeding MR, et al. "Efficacy and long-term safety of a dengue vaccine in regions of endemic disease." *N Engl J Med* (2015): 1195-1206.
2. World Health Organization. *Background paper on Dengue Vaccines prepared by SAGE working group on Dengue vaccines and WHO secretariat*. Geneva, 17 March 2016. Accessed at: http://www.who.int/immunization/sage/meetings/2016/april/1_Background_Paper_Dengue_Vaccines_2016_03_17.pdf
3. Philippine Society of Microbiology and Infectious Diseases. *PSMID Statement on Dengue Vaccination in Adults*. 25 October 2016. Accessed at <<https://www.pcp.org.ph/index.php/latest-news-announcements/658-psmid-statement-on-dengue-vaccination-in-adults>>.
4. Sanofi. *Sanofi updates information on dengue vaccine*. 29 November 2017. Accessed at <<http://mediaroom.sanofi.com/sanofi-updates-information-on-dengue-vaccine/>>.
5. WHO. *Updated Questions and Answers related to information presented in the Sanofi Pasteur press release on 30 November 2017 with regards to the dengue vaccine Dengavaxia*. 30 November 2017. <http://www.who.int/immunization/diseases/dengue/q_and_a_dengue_vaccine_dengavaxia/en/>.
6. Villar L, Dayan GH, Arredondo-García JL, et al. "Efficacy of a tetravalent dengue vaccine in children in Latin America." *N Eng J Med* 372.Supplement (2015):113-23.
7. WHO. (2009). *Dengue: Guidelines for diagnosis, treatment, prevention and control*.